

The final fate of instability of Reissner-Nordström-anti-de Sitter black holes by charged complex scalar fields

Kengo Maeda* and Shunsuke Fujii†

Faculty of Engineering, Shibaura Institute of Technology, Saitama, 330-8570, Japan

Jun-ichirou Koga‡

Advanced Research Institute for Science and Engineering, Waseda University, Shinjuku, Tokyo 169-8555

(Dated: May 24, 2010)

We investigate instability of 4-dimensional Reissner-Nordström-anti-de Sitter (RN-AdS₄) black holes with various topologies by charged scalar field perturbations. We numerically find that the RN-AdS₄ black holes become unstable against the linear perturbations below a critical temperature. It is analytically shown that charge extraction from the black holes occurs during the unstable evolution. To explore the end state of the instability, we perturbatively construct static black hole solutions with the scalar hair near the critical temperature. It is numerically found that the entropy of the hairy black hole is always larger than the one of the unstable RN-AdS₄ black hole in the microcanonical ensemble. Our results support the speculation that the black hole with charged scalar hair always appears as the final fate of the instability of the RN-AdS₄ black hole.

PACS numbers: 04.70.Bw, 04.70.Dy, 11.25.Tq

I. INTRODUCTION

Motivated by a holographic model of superconductors, there has been renewal of interest in the Einstein-Maxwell-scalar system in asymptotically anti-de Sitter spacetime. The model is constructed by a gravitational theory of a charged complex scalar field coupled to the Maxwell field (for review, see Refs. [1–3]) via AdS/CFT (anti-de Sitter/conformal field theory) duality [4]. According to the AdS/CFT dictionary, a condensate in the holographic superconductor is described by a hairy black hole dressed with the charged scalar field. It was numerically found that the hairy black hole exists below a critical temperature T_c in the plane-symmetric spacetime and the phase transition is second-order [5].

For a gravity system minimally coupled with a neutral scalar field, it is well known that the scalar field perturbation with mass satisfying Breitenlohner-Freedman (BF) bound does not cause any instabilities of AdS spacetime [6]. So, even though a black hole solution with the scalar hair exists in asymptotically AdS spacetime [7], Schwarzschild-AdS (Sch-AdS) spacetime should be stable against the scalar field perturbations if BF bound is satisfied. In 4 and 5-dimensional $N = 8$ gauged supergravities, this speculation was supported by the numerical calculation that the entropy of the Sch-AdS black hole is larger than the one of the hairy black hole in the microcanonical ensemble [8]. In this paper we investigate how this picture is modified in the gravity system minimally coupled with a charged scalar field with mass satisfying BF bound. As discussed in Ref. [9], RN-AdS black hole

with low temperature may be unstable against the perturbation of the charged scalar field since the effective mass of the scalar field is negative near the horizon. So, we naively expect that a black hole with the scalar hair appears as the final fate of the instability even though BF bound is satisfied.

In AdS spacetime, the Dirichlet boundary condition is usually imposed at spatial infinity when we consider the dynamical evolution. Under the boundary condition, energy flux and charge current do not enter through the infinity, and therefore, the total mass and charge of the spacetime are conserved during the dynamical evolution. So, if RN-AdS₄ black hole evolves to the hairy black hole as the end state of the instability, the entropy of the hairy black hole should be larger than the one of RN-AdS₄ black hole in the microcanonical ensemble due to the second law of black hole.

To explore the above speculation we first investigate instability of RN-AdS₄ black holes with various topologies of horizon by charged complex scalar fields with various mass satisfying BF bound. Second, we numerically show that the entropy of the hairy black hole is always larger than the one of RN-AdS₄ black hole whenever the RN-AdS₄ black hole is unstable. An analytic calculation also shows that charge is extracted from the unstable black hole via the charged scalar field.

The plan of our paper is as follows: In Sec. II, we investigate the lowest quasi-normal (QN) frequencies of RN-AdS₄ black holes both numerically and analytically near the second order phase transition. In Sec. III, we construct the hairy black hole solutions by solving the field equations perturbatively near the phase transition. The entropy of the hairy black hole is numerically calculated by using the first law of black hole. Conclusion and discussions are devoted to Sec. IV.

*Electronic address: maeda302@sic.shibaura-it.ac.jp

†Electronic address: fujii@sic.shibaura-it.ac.jp

‡Electronic address: koga@gravity.phys.waseda.ac.jp

II. QUASI-NORMAL FREQUENCIES NEAR THE SECOND ORDER PHASE TRANSITION

We consider the Lagrangian density of Einstein-Maxwell-scalar system in 4-dimensional asymptotically anti-de Sitter (AdS₄) spacetime:

$$\begin{aligned}\mathcal{L} &= R + \frac{6}{L^2} - \frac{F^{ab}F_{ab}}{4} - m^2|\psi|^2 - |D\psi|^2, \\ D_a &= \nabla_a - iqA_a, \quad F_{ab} := \partial_a A_b - \partial_b A_a\end{aligned}\quad (2.1)$$

where L , m , and q are the AdS radius, the mass, and the charge of the complex scalar field, respectively. The equations of motion are given by

$$D_a D^a \psi = m^2 \psi, \quad (2.2a)$$

$$\nabla_a F_b{}^a = j_b, \quad j_a = iq[(D_a \psi)^\dagger \psi - (D_a \psi) \psi^\dagger], \quad (2.2b)$$

$$\begin{aligned}G_{ab} &= \frac{3g_{ab}}{L^2} + \frac{1}{2}F_{ac}F_b{}^c - \frac{g_{ab}}{8}F^2 - \frac{m^2}{2}|\psi|^2 g_{ab} \\ &\quad - \frac{g_{ab}}{2}|D\psi|^2 + \frac{1}{2}[D_a \psi (D_b \psi)^\dagger + (D_a \psi)^\dagger D_b \psi].\end{aligned}\quad (2.2c)$$

The simple solution of the equations are given by Reissner-Nordström (RN)-AdS₄ black hole solutions with various curvature of horizon. The metric is given by

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{f(r)} + r^2 d\Omega_2^2, \quad (2.3a)$$

$$\begin{aligned}f(r) &= k - \frac{2M}{r} + \frac{\rho^2}{4r^2} + \frac{r^2}{L^2} \\ &= \frac{1}{L^2 r^2}(r - r_+)(r - r_-)(r^2 + ar + b),\end{aligned}\quad (2.3b)$$

$$\begin{aligned}a &:= r_+ + r_-, \quad b := kL^2 + (r_+ + r_-)^2 - r_+ r_-, \\ T &= \frac{(r_+ - r_-)(r_+^2 + ar_+ + b)}{4\pi L^2 r_+^2},\end{aligned}\quad (2.3c)$$

where $d\Omega_2^2$, r_+ (r_-), and T are the metric of 2-dimensional Einstein space with constant scalar curvature $2k$ ($k = 0, \pm 1$), the outer (inner) horizon radius, and the Hawking temperature, respectively. The mass (charge) parameter M (ρ) corresponds to the total mass (charge) for $k = 1$ and the mass (charge) density for $k = 0, -1$, respectively. These two parameters are rewritten in terms of $r_\pm$, L , and k as

$$\rho = 2\sqrt{\frac{r_+ r_- (kL^2 + (r_+ + r_-)^2 - r_+ r_-)}{L^2}}, \quad (2.4a)$$

$$M = \frac{r_+ + r_-}{2L^2}(kL^2 + r_+^2 + r_-^2), \quad (2.4b)$$

and the gauge potential A_μ is given by

$$A_\mu dx^\mu = \Phi(r)dt = \rho \left(\frac{1}{r_+} - \frac{1}{r} \right) dt. \quad (2.5)$$

We obtain the lowest quasi-normal (QN) frequencies of RN-AdS₄ black holes by considering perturbation of the

solutions. Since $\psi = 0$ on the RN-AdS₄ solutions, the scalar perturbation decouples from the perturbations of the other fields. So, we solve Eq. (2.2a) for the fixed metric (2.3a) and the gauge potential (2.5). Near the horizon, ψ behaves as

$$\psi_\mp \sim e^{-i\omega t}(1-u)^{\mp\gamma\omega i}, \quad \gamma = \frac{L^2 r_+^2}{(r_+ - r_-)(r_+^2 + ar_+ + b)}, \quad (2.6)$$

where u is defined by $u := r_+/r$. From the retarded condition, we must impose the “ingoing wave” boundary condition: $\psi \sim \psi_-$ on the horizon. Defining a new variable Ψ as

$$\psi(t, u) = e^{-i\omega t}(1-u)^{-\gamma\omega i}\Psi(u), \quad (2.7)$$

Eq. (2.2a) is written as

$$\begin{aligned}\Psi''(u) &+ \left(\frac{2\gamma\omega i}{1-u} + \frac{f'(u)}{f(u)} \right) \Psi'(u) \\ &+ \left[\frac{\gamma\omega i}{1-u} \left(\frac{f'(u)}{f(u)} + \frac{\gamma\omega i + 1}{1-u} \right) \right. \\ &\quad \left. + \frac{r_+^2}{u^4 f^2(u)}(\omega + qA_t)^2 - \frac{r_+^2 m^2}{u^4 f(u)} \right] \Psi(u) = 0, \\ f(u) &= \frac{r_+^2}{L^2 u^2}(1-u) \left(1 - \frac{r_-}{r_+} u \right) \left(1 + \frac{a}{r_+} u + \frac{b}{r_+^2} u^2 \right),\end{aligned}\quad (2.8)$$

where a prime denotes the derivative with respect to u . In terms of this variable Ψ , the “ingoing wave” boundary condition is described by the regularity condition:

$$\Psi(u) \text{ is regular at } u = 1. \quad (2.9)$$

Near the spatial infinity, Ψ behaves as

$$\Psi(u) = c_1 u^{\Delta_+} + c_2 u^{\Delta_-}, \quad \Delta_\pm = \frac{3}{2} \pm \sqrt{9/4 + L^2 m^2}, \quad (2.10)$$

where Breitenlohner-Freedman (BF) bound [6] is represented by $m^2 \geq -9/4L^2$. Since we are concerned with the time evolution under which both the mass (or mass density) M and the charge (or charge density) ρ are conserved, we shall impose Dirichlet boundary condition at infinity¹:

$$\Psi(u) \sim u^{\Delta_+}. \quad (2.11)$$

According to the critical slowing down phenomenon, the lowest QN frequencies ω should approach zero toward

¹ When $-9/4L^2 \leq m^2 < -5/4L^2$, there are two normalizable modes. So, imposing $c_1 = 0$ instead of $c_2 = 0$ is also possible as the boundary condition. It is well known, however, that the boundary condition in which neither c_1 nor c_2 vanishes is unphysical since the spacetime is unstable [12].

the second order phase transition $T = T_c$ [10, 11]. This implies that one can expand the solution of Eq. (2.8) as

$$\Psi(u) = \Psi_0(u) + \omega \Psi_1(u) + O(\omega^2) \quad (2.12)$$

near the phase transition, where $\Psi_0(u)$ is the marginally stable solution. Denoting the deviations from the phase transition as

$$\begin{aligned} r_+ &= r_c + \delta r_+, \\ \gamma &= \gamma_c + \delta \gamma, \\ f(u) &= f_c(u) + \delta f(u), \\ \Phi(u) &= \Phi_c(u) + \delta \Phi(u), \end{aligned} \quad (2.13)$$

we write down the equations for $\Psi_0(u)$ and $\Psi_1(u)$ as

$$\mathcal{L}\Psi_0(u) = 0, \quad (2.14)$$

$$\mathcal{L}(\omega \Psi_1) = j_1(u), \quad (2.15)$$

where the differential operator $\mathcal{L}$ and the source term j_1 are defined by

$$\mathcal{L} := \frac{d^2}{du^2} + \frac{f'_c(u)}{f_c(u)} \frac{d}{du} + \frac{r_c^2}{u^4 f_c(u)} \left(\frac{q^2 \Phi_c^2(u)}{f_c(u)} - m^2 \right), \quad (2.16)$$

$$\begin{aligned} \text{Re}[j_1] &:= \frac{2\gamma_c \omega_I}{1-u} \Psi'_0(u) - \delta \left(\frac{f'(u)}{f(u)} \right) \Psi'_0(u) \\ &+ \left[\frac{\gamma_c \omega_I}{1-u} \left(\frac{f'_c(u)}{f_c(u)} + \frac{1}{1-u} \right) - \frac{2q^2 r_c \Phi_c^2(u)}{u^4 f_c(u)} \delta \left(\frac{r_+}{f(u)} \right) \right. \\ &- \frac{2q \omega_R r_c^2}{u^4 f_c^2(u)} \Phi_c(u) - \frac{2q^2 r_c^2 \Phi_c(u) \delta \Phi(u)}{u^4 f_c^2(u)} \\ &\left. + \frac{2m^2 r_c \delta r_+}{u^4 f_c(u)} - \frac{m^2 r_c^2 \delta f(u)}{u^4 f_c^2(u)} \right] \Psi_0(u), \end{aligned} \quad (2.17a)$$

$$\begin{aligned} \text{Im}[j_1] &:= -\frac{\gamma_c \omega_R}{1-u} \left[2\Psi'_0(u) + \left(\frac{f'_c(u)}{f_c(u)} + \frac{1}{1-u} \right) \Psi_0(u) \right] \\ &- \frac{2q \omega_I r_c^2}{u^4 f_c^2(u)} \Phi_c(u) \Psi_0(u) \end{aligned} \quad (2.17b)$$

for $\omega = \omega_R + i\omega_I$. To derive Eq. (2.17), we assumed that Ψ_0 is a real function of u without loss of generality.

To derive ω from Eq. (2.15), let us define the inner product as

$$\langle \psi_a | \psi_b \rangle := \int_0^1 \psi_a^\dagger(u) \psi_b(u) du \quad (2.18)$$

for any functions ψ_a satisfying both boundary conditions (2.9) and (2.11). Under these boundary conditions, the differential operator $\hat{\mathcal{L}} := f_c(u)\mathcal{L}$ is shown to be an Hermite operator and by Eq. (2.14) we obtain

$$\langle \hat{j}_1 | \Psi_0 \rangle = \langle \hat{\mathcal{L}} \Psi_1 | \Psi_0 \rangle = \langle \Psi_1 | \hat{\mathcal{L}} \Psi_0 \rangle = 0, \quad (2.19)$$

where $\hat{j}_1(u) := f_c(u)j_1(u)/\omega$. The real and imaginary parts of Eq. (2.19) yield the lowest QN frequencies ω as

$$\omega_R = -\frac{2q r_c \omega_I}{\Psi_0^2(1)} \int_0^1 \frac{\Phi_c(u) \Psi_0^2(u)}{u^4 f_c(u)} du, \quad (2.20a)$$

$$\begin{aligned} \omega_I r_c &\left[\Psi_0^2(1) + \frac{4q^2 r_c^2}{\Psi_0^2(1)} \left(\int_0^1 \frac{\Phi_c(u) \Psi_0^2(u)}{u^4 f_c(u)} du \right)^2 \right] \\ &= \frac{1}{2} \int_0^1 \delta \left(\frac{f'(u)}{f(u)} \right) f_c(u) \{ \Psi_0^2(u) \}' du \\ &+ 2q^2 r_c \int_0^1 \frac{\Psi_0^2(u)}{u^4} \delta \left(\frac{r_+}{f(u)} \right) \Phi_c^2(u) du \\ &+ \int_0^1 \frac{\Psi_0^2(u)}{u^4 f_c(u)} \left[q^2 r_c^2 \delta(\Phi^2(u)) - m^2 f_c^2(u) \delta \left(\frac{r_+^2}{f(u)} \right) \right] du. \end{aligned} \quad (2.20b)$$

It is noteworthy that ω is independent of the absolute amplitude of $\Psi_0(1)$ and it is determined by solving only the leading order equation (2.14).

Hereafter, we shall fix L to $L = 1$. Then, from Eqs. (2.3c) and (2.4), $r_\pm$ and T are functions of M for given k and ρ . So, for the sake of later convenience, we shall define the deviation parameter from the second order phase transition as

$$\epsilon := \frac{M_c - M}{M_c}, \quad (2.21)$$

where M_c is the mass at the critical temperature $T = T_c$. Since M is locally a smooth function of k , ρ , and T , ϵ approaches zero in the limit $T \rightarrow T_c$. From Eqs. (2.4) $\delta r_\pm$ is related to ϵ as

$$\delta r_- = -\frac{r_c^*(kL^2 + 3r_c^2 + 2r_c r_c^* + r_c^{*2})}{r_c(kL^2 + 3r_c^{*2} + 2r_c r_c^* + r_c^2)} \delta r_+ \quad (2.22a)$$

and

$$\delta r_+ = -\frac{(r_c + r_c^*)(kL^2 + r_c^2 + r_c^{*2})\epsilon}{(kL^2 + 3r_c^2 + 2r_c r_c^* + r_c^{*2})(1 - r_c^*/r_c)}, \quad (2.22b)$$

where r_c^* is the inner horizon radius r_- at $T = T_c$. To derive Eq. (2.22), we impose the condition that ρ is constant along the deviation.

For a generic value of M both of the two boundary conditions (2.9) and (2.11) are not necessarily satisfied, as Eq. (2.14) is the second order linear differential equation. By using Mathematica, we numerically scan through possible values of M until the two boundary conditions are satisfied and find M_c for given m , q , k , and ρ . Then, representing δf , $\delta \Phi$, and δr_+ by ϵ via Eqs. (2.22) and integrating Eqs. (2.20) numerically, we obtain ω_I .

Since all the deviations are proportional to ϵ , ω is also proportional to ϵ , in agreement with the critical slowing down [10, 11]. We find that ω_I becomes positive (negative) for $\epsilon > 0$ (< 0), independent of k , q , and m^2 . In Fig. 1 we plot q -dependence of the dimensionless quantity




FIG. 1: (color online). The q -dependence of $-r_c^2 \omega_I / \delta r_+$ is shown for various k in the case of $m^2 = -2/L^2$ and $\rho = 2$. The star, box, and diamond represent $k = 1$, $k = 0$, and $k = -1$, respectively.




FIG. 2: (color online). The q -dependence of $-r_c^2 \omega_I / \delta r_+$ is shown for several m^2 in the case of $k = 0$ and $\rho = 60$. The star, box, and diamond represent $m^2 = -2/L^2$, $m^2 = 0$, and $m^2 = 7/4 L^2$, respectively. At the left edge of each plot, the RN-AdS₄ black holes become almost extremal.

$-r_c^2 \omega_I / \delta r_+$ for various k in the case of $m^2 = -2/L^2$ and $\rho = 2$. In Fig. 2, we plot q -dependence of $-r_c^2 \omega_I / \delta r_+$ for several m^2 in the case of $\rho = 60$. These figures show that RN-AdS₄ black holes become unstable against the linear perturbation of the complex scalar field when $T < T_c$ ($\epsilon > 0$),² as $\epsilon / \delta r_+ < 0$, independent of k and m^2 .

The charge current through the horizon is calculated by contracting j_a in Eq. (2.2b) with the killing vector $k^a = (\partial_t)^a$, which is null on the horizon. By Eq. (2.20a),

² We assumed that $\epsilon > 0$ for $T < T_c$. This corresponds to the assumption that the specific heat of the RN-AdS₄ black hole is positive at $T = T_c$.

we obtain

$$j_a k^a|_{r=r_c} = 4q^2 r_c \omega_I e^{2\omega_I t} \int_0^1 \frac{\Phi_c(u) \Psi_0^2(u)}{u^4 f_c(u)} du, \quad (2.23)$$

which shows that $j_a k^a|_{r=r_c}$ is positive when $\omega_I > 0$. This implies that charge extraction occurs from the RN-AdS₄ black hole when it is unstable.

III. THE ENTROPY FOR HAIRLY BLACK HOLES

In this section, we investigate the final fate of the instabilities found in Sec. II by calculating the entropy of the hairy black holes. We start by constructing the static hairy black hole solutions near the phase transition. Near the phase transition, the equations of motion (2.2) can be perturbatively solved since the scalar field ψ is very small. To solve the perturbed equations easily, we adopt a coordinate system $\{T, y\}$ where the location of the horizon is set to be $y = 1$. So, we shall take the following ansatz for the metric and the gauge field:

$$ds^2 = -g(y)dT^2 + \frac{dy^2}{g(y)} + R^2(y)d\Omega_2^2, \quad (3.1a)$$

$$A_\mu dx^\mu = \phi(y)dT. \quad (3.1b)$$

Assuming that ψ is real without loss of generality, the scalar and Maxwell equations (2.2a) and (2.2b) are reduced to the two coupled differential equations:

$$g\psi'' + \left(\frac{2gR'}{R} + g'\right)\psi' + \frac{q^2\phi^2\psi}{g} = m^2\psi, \quad (3.2a)$$

$$\phi'' + \frac{2R'}{R}\phi' = \frac{2q^2\phi\psi^2}{g}, \quad (3.2b)$$

where a prime denotes the derivative with respect to y . The metric functions g and R are obtained by solving the Einstein equations (2.2c):

$$\frac{R''}{R} = -\frac{q^2\phi^2\psi^2}{2g^2} - \frac{\psi'^2}{2}, \quad (3.3a)$$

$$\begin{aligned} \frac{R'}{R}g' &= \frac{k}{R^2} - \frac{gR'^2}{R^2} + \frac{3}{L^2} - \frac{1}{4}\phi'^2 - \frac{1}{2}m^2\psi^2 \\ &+ \frac{1}{2}g\psi'^2 + \frac{q^2}{2g}\phi^2\psi^2. \end{aligned} \quad (3.3b)$$

As shown in Refs. [13, 14], we can expand the functions ψ , g , R , ϕ , and temperature T near the second order phase transition as

$$\begin{aligned} \psi(y) &= \epsilon^{1/2}\psi_1(y) + \epsilon^{3/2}\psi_2(y) + \dots, \\ g(y) &= g_c(y) + \epsilon g_1(y) + \dots, \\ R(y) &= r_c y + \epsilon R_1(y) + \dots, \\ \phi(y) &= \phi_c(y) + \epsilon \phi_1(y) + \dots, \\ T(\epsilon) &= T_c + \epsilon T_1 + \dots. \end{aligned} \quad (3.4)$$

The zeroth order solutions $g_c(y)$ and $\phi_c(y)$ are simply given by the coordinate transformation:

$$r = r_c y, \quad t = T/r_c \quad (3.5)$$

of the RN-AdS₄ black hole solution (2.3) as

$$g_c(y) = \frac{1}{L^2 y^2} (y-1)(y-r_c^*/r_c) \left(y^2 + \frac{a}{r_c} y + \frac{b}{r_c^2} \right), \quad (3.6a)$$

$$\phi_c(y) = \frac{\rho}{r_c^2} \left(1 - \frac{1}{y} \right). \quad (3.6b)$$

Then, $\psi_1(y)$, $\phi_1(y)$, $R_1(y)$, and $g_1(y)$ obey the following linear differential equations:

$$\frac{(y^2 g_c \psi_1')'}{y^2} = \mathcal{M}^2 \psi_1, \quad (3.7)$$

$$\phi_1'' + \frac{2}{r_c} \left(\frac{R_1'}{y} - \frac{R_1}{y^2} \right) \phi_1' + \frac{2}{y} \phi_1' = \frac{2q^2 \phi_c \psi_1^2}{g_c}, \quad (3.8)$$

$$R_1'' = -\frac{r_c y}{2} \left(\frac{q^2 \phi_c^2 \psi_1^2}{g_c^2} + \psi_1'^2 \right), \quad (3.9)$$

$$\begin{aligned} \frac{1}{y} g_1' + \frac{g_1}{y^2} + \frac{1}{r_c} \left(g_c' + \frac{2g_c}{y} \right) \left(\frac{R_1'}{y} - \frac{R_1}{y^2} \right) g_c' + \frac{2kR_1}{r_c^3 y^3} \\ = -\frac{1}{2} \phi_c' \phi_1' - \frac{m^2}{2} \psi_1^2 + \frac{1}{2} g_c \psi_1'^2 + \frac{q^2}{2g_c} \phi_c^2 \psi_1^2, \end{aligned} \quad (3.10)$$

where $\mathcal{M}$ is the effective mass defined by

$$\mathcal{M}^2(y) := m^2 - q^2 \phi_c^2 / g_c \quad (3.11)$$

which can be negative [9].

The solutions of Eqs. (3.8) and (3.9) are formally given by

$$\phi_1' = \frac{2}{y^2} \int_{\infty}^y \left[\frac{q^2 \phi_c \psi_1^2 z^2}{g_c} + \frac{\phi_c'}{r_c} (R_1 - z R_1') \right] dz + \frac{c_3}{y^2}, \quad (3.12a)$$

$$R_1(y) = \mathcal{R}(y) + c_1 y + c_2, \quad (3.12b)$$

$$\mathcal{R}(y) := -\frac{r_c}{2} \int_{\infty}^y \int_{\infty}^z \left(\frac{q^2 \phi_c^2(w) \psi_1^2(w)}{g_c^2(w)} + \psi_1'^2(w) \right) w dw dz, \quad (3.12c)$$

where c_1 , c_2 , and c_3 are integral constants. Note that $\psi_1(y)$ is given by $\Psi_0(1/y)$, as Eq. (3.7) is equivalent to Eq. (2.14) by setting $y = 1/u$. So, all solutions are determined by Ψ_0 except the integral constants.

Since ψ_1 rapidly decays as $\psi = O(y^{-\lambda})$ ($\lambda > 3/2$) near infinity, the first term of Eq. (3.12a) decays as $O(y^{-3})$,

which does not contribute to the charge (or charge density) ρ_h of the hairy black hole solutions. ρ_h is given by

$$\rho_h = \rho \left(1 + \frac{2c_1 \epsilon}{r_c} + \frac{r_c^2 c_3 \epsilon}{\rho} \right) \quad (3.13)$$

up to the order $O(\epsilon)$. So, equating $\rho_h = \rho$, we obtain the constant c_3 as

$$c_3 = -\frac{2c_1 \rho}{r_c^3}. \quad (3.14)$$

The formal solution of Eq. (3.10) is given by

$$\begin{aligned} g_1 = \frac{1}{r_c y} \int_1^y (R_1 - R_1' y) \left(g_c' + \frac{2g_c}{y} \right) dy - \frac{2}{y} \int_1^y \frac{kR_1}{r_c^3 y} dy \\ - \frac{1}{2y} \int_1^y y^2 \phi_c' \phi_1' dy - \frac{m^2}{2y} \int_1^y y^2 \psi_1^2 dy \\ + \frac{1}{2y} \int_1^y y^2 g_c \psi_1'^2 dy + \frac{q^2}{2y} \int_1^y \frac{y^2 \phi_c^2 \psi_1^2}{g_c} dy. \end{aligned} \quad (3.15)$$

To read off the mass (or mass density) M from the metric (3.1a), we need to take the coordinate transformation $T \rightarrow t$:

$$T = t(r_c + c_1 \epsilon) \quad (3.16)$$

so that the leading term of the metric (3.1a) near the spatial infinity $y \rightarrow \infty$ coincides with the asymptotic form of the metric (2.3) of RN-AdS₄ black holes:

$$ds^2 = -\frac{R^2}{L^2} dt^2 + \frac{L^2 dR^2}{R^2} + R^2 d\Omega_2^2. \quad (3.17)$$

Substituting Eqs. (3.12) and (3.14) into Eq. (3.15), we find that the asymptotic fall-off of $-g_{tt}$ component behaves as

$$\begin{aligned} -g_{tt} = f_c(R) + \frac{2\epsilon \chi(\psi_1)}{R} - \frac{4\pi(c_1 + c_2)r_c T_c}{R} \epsilon \\ + \dots + O(\epsilon^2) \\ = \frac{R^2}{L^2} + k - \frac{2M}{R} + \dots + O(\epsilon^2). \end{aligned} \quad (3.18)$$

Here, χ is a complicated function of ψ_1 and it vanishes when $\psi_1 = 0$. By Eqs. (2.21) and (3.18), the combination of the constants $c_1 + c_2$ is rewritten by χ and M_c as

$$c_1 + c_2 = \frac{\chi(\psi_1) - M_c}{2\pi T_c r_c}. \quad (3.19)$$

Note that RN-AdS₄ black hole solutions with mass (or mass density) $M = M_c(1 - \epsilon)$ are automatically derived by setting $\psi_1 = \chi(\psi_1) = 0$.

We calculate the entropy (or entropy density) S of the hairy black holes with the same mass (or mass density) M by using the first law (A18) for a fixed charge (or charge density) ρ near the critical temperature T_c :

$$\frac{\delta S}{\delta M}(\epsilon) = \frac{1}{T(\epsilon)} = \frac{1}{T_c} - \frac{T_1}{T_c^2} \epsilon + O(\epsilon^2). \quad (3.20)$$

Integrating this differential equation under the relation $\delta M = -M_c \delta \epsilon$, we obtain

$$\begin{aligned} S &= S_c + S_1 \epsilon + S_2 \epsilon^2 + \dots \\ &= S_c - \frac{M_c \epsilon}{T_c} + \frac{M_c T_1 \epsilon^2}{2T_c^2} + O(\epsilon^3). \end{aligned} \quad (3.21)$$

This means that the difference of the entropy (or entropy density) between RN-AdS₄ black hole and the hairy black hole appears at $O(\epsilon^2)$:

$$\begin{aligned} \Delta S &= \frac{\epsilon^2 M_c}{2T_c^2} \Delta T_1 = S_{\text{hairlyBH}} - S_{\text{RN-AdS}_4\text{BH}} \\ &\propto R^2(1)_{\text{hairlyBH}} - R^2(1)_{\text{RN-AdS}_4\text{BH}} = O(\epsilon^2), \end{aligned} \quad (3.22)$$

and then $R_1(1)|_{\text{hairlyBH}} = R_1(1)|_{\text{RN-AdS}_4\text{BH}}$. Hence, by Eqs. (3.12b) and (3.19), $c_1 + c_2$ is rewritten by $\mathcal{R}$ as

$$c_1 + c_2 = -\mathcal{R}(1) - \frac{M_c}{2\pi T_c r_c}. \quad (3.23)$$

As a check of numerical calculation below, we verified that the r.h.s. of Eq. (3.19) agrees with the r.h.s. of Eq. (3.23) within the numerical error.

We obtain the Hawking temperature T from the regularity of the Euclidean solution analytically continued by $t \rightarrow i\tau$ as

$$T(\epsilon) = T_c + \frac{r_c}{4\pi} \left(g'_1(1) + \frac{c_1}{r_c} g'_c(1) \right) \epsilon + O(\epsilon^2), \quad (3.24)$$

where the critical temperature T_c is given by $T_c = r_c g'_c(1)/4\pi$. By using Eqs. (3.12), (3.15), and (3.23), T_1 in Eq. (3.4) is expressed by

$$\begin{aligned} T_1 &= \frac{r_c}{4\pi} \left(g'_1(1) + \frac{c_1}{r_c} g'_c(1) \right) \\ &= \frac{1}{4\pi} \left[-\frac{r_c g'_c(1)}{2} \int_1^\infty \left(\frac{q^2 \phi_c^2(y) \psi_1^2(y)}{g_c^2(y)} + \psi_1'^2(y) \right) y dy \right. \\ &\quad \left. + \frac{\rho}{r_c} \int_1^\infty \frac{q^2 \phi_c(y) \psi_1^2(y) y^2}{g_c(y)} dy - \frac{m^2}{2} r_c \psi_1^2(1) \right] \\ &\quad - \frac{M_c}{8\pi^2 T_c r_c^3} \left(4\pi T_c r_c + \frac{\rho^2}{r_c^2} \right). \end{aligned} \quad (3.25)$$

The first term in the second equality corresponds to the difference of T_1 between the RN-AdS₄ black hole and the hairy one. Eliminating $\psi_1'^2$ term by Eq. (3.7) and using integration by parts, we finally obtain a useful expression for the difference $\Delta T_1 := T_1|_{\text{hairlyBH}} - T_1|_{\text{RN-AdS}}$ as

$$\begin{aligned} \frac{\Delta T_1}{T_c} &= \frac{\psi_1^2(1)}{4} + \frac{1}{2} \int_1^\infty (m^2 \psi_1 - g'_c \psi_1') \frac{\psi_1}{g_c} y dy \\ &\quad + \frac{1}{L^2 g'_c(1)} \int_1^\infty \frac{(y-1)y}{g_c} \left(\frac{kL^2}{r_c^2} + 3 + y(2+y) \right) \frac{q^2 \phi_c^2 \psi_1^2}{g_c} dy. \end{aligned} \quad (3.26)$$

Note that the first integral does not diverge according to the boundary condition at the horizon; $g'_c(1)\psi_1'(1) = m^2\psi_1(1)$.




FIG. 3: (color online). The q -dependence of ΔS_2 for the solution of Eq. (3.7) normalized as $\psi_1(1) = 1$ in the case of $m^2 = -2/L^2$, $\rho = 2$. The star, box, and diamond represent $k = 1$, $k = 0$, and $k = -1$, respectively.

By demanding that the critical temperature in Eq. (2.3c) is non-negative, we obtain an inequality $kL^2/r_c^2 + 6 > 0$, which guarantees the positivity of the second integral. So, by Eq. (3.21), we observe that the negative part of the effective mass (3.11), $q^2 \phi_c^2/g_c$, increases the entropy of the hairy black hole at the second order $O(\epsilon^2)$.

If the hairy black hole appears as the final fate of the instabilities found in Sec. II, the entropy of the hairy black hole should be larger than the one of the RN-AdS₄ black hole according to the second law of the black hole. To check this, we numerically calculate the entropy difference between the hairy black hole and the RN-AdS₄ black hole:

$$\Delta S_2 := S_2|_{\text{hairlyBH}} - S_2|_{\text{RN-AdS}_4\text{BH}} \quad (3.27)$$

by integrating Eq. (3.26) for various values of m^2 and k . Since we are only interested in the sign of ΔS_2 , we numerically calculate ΔS_2 for the solution of Eq. (3.7) by normalizing as $\psi_1(1) = 1$.

In Fig. 3, we plot ΔS_2 for various q and k in the case of $m^2 = -2/L^2$ and $\rho = 2$. Fig. 4 shows the q -dependence of $\text{Log}_{10}(\Delta S_2)$ for various m^2 in the case of $k = 0$ and $\rho = 60$. Both figures indicate that ΔS_2 is positive, independent of the mass m^2 , charge q , and curvature k .

We also give a further evidence that the hairy black hole constructed above appears as the end state of the instabilities found in Sec. II by calculating charge density $\tilde{\rho}$ outside the black hole horizon. $\tilde{\rho}$ on $T = \text{const.}$ hypersurface Σ is calculated at $O(\epsilon)$ as

$$\tilde{\rho} = j_a n^a = 2\epsilon \frac{q^2 \phi_c(y)}{\sqrt{g_c(y)}} \psi_1^2 = \frac{2\epsilon q^2 \rho \left(1 - \frac{1}{y}\right)}{r_c^2 \sqrt{g_c(y)}} \psi_1^2(y) > 0, \quad (3.28)$$

where n^a is the past directed unit normal vector on Σ . So, positive charge is distributed outside the hairy black




FIG. 4: (color online). The q -dependence of $\text{Log}_{10}(\Delta S_2)$ for the solution of Eq. (3.7) normalized as $\psi_1(1) = 1$ for various m^2 in the case of $k = 0$, $\rho = 60$. The star, box, and diamond represent $m^2 = -2/L^2$, $m^2 = 0$, and $m^2 = 7/4 L^2$, respectively. At the left edge of each plot, the RN-AdS₄ black holes become almost extremal.

hole. This agrees with the result (2.23), where the positive charge is shown to be extracted from the horizon via the complex scalar field when RN-AdS₄ black hole becomes unstable.

IV. CONCLUSION AND DISCUSSIONS

We have investigated instability of RN-AdS₄ black holes with various topologies against perturbation of charged scalar fields with various masses near the second order phase transition. Near the phase transition where the scalar field begins to condense, all fields are expected to evolve very slowly according to the critical slowing down [10, 11]. So, we obtained a useful expression for the lowest QN frequencies of the RN-AdS₄ black holes by solving the scalar field equation (2.8) perturbatively around the the phase transition. It is numerically shown that RN-AdS₄ black holes become unstable against the perturbations below the critical temperature T_c . By Eq. (2.23), we found that the charge extraction occurs from the RN-AdS₄ black holes during the unstable evolution.

To see the final fate of the instabilities, we have also constructed the black hole solutions with the scalar hair by solving the field equations (2.2) perturbatively around the the phase transition. We found that the negative part of the effective mass $\mathcal{M}^2$ of the scalar field (3.11) increases the entropy. The numerical calculations indicate that the entropy of the hairy black holes is always larger than the one of RN-AdS₄ black holes in the microcanonical ensemble whenever RN-AdS₄ black holes are unstable. As indicated in Eqs. (3.28) and (2.23), the positive charge density outside the hairy black hole is extracted from the RN-AdS₄ black hole. These facts suggest that the hairy black holes are the final fate of the instabilities of RN-AdS₄ black holes, independent of the topology of

the horizon and mass of the scalar field. So, this is a universal feature of the charged complex scalar system.

Since the charge is extracted from the RN-AdS₄ black holes, the instability against the complex scalar field is very similar to the superradiance phenomenon where energy is extracted from a rotating black hole. The unstable modes inspired by superradiance were recently found in Kerr-AdS black hole [15] only when the Einstein universe at the AdS boundary rotates faster than the speed of light. This is due to the fact that a global timelike killing vector field does not exist in the rapidly rotating case [16, 17]. Refs. [16, 17] also claim that there is no unstable mode for any linear perturbation equation when spacetime possesses a global timelike killing vector field $k^a = (\partial_t)^a$ and the energy current $J^a := -T_{ab}k^b$ satisfies the conservation law $\nabla_a J^a = 0$ and the dominant energy condition, i.e., $J^a V_a \leq 0$ for any future-directed timelike vector V^a .

At first sight, this seems to contradict the unstable modes found in this paper, as they satisfy the dominant energy condition for $m^2 \geq 0$ and RN-AdS₄ spacetime possesses the global timelike killing vector field k^a . However, the energy current J^a of the scalar field is not conserved for the *fixed* gauge potential A_a in Eq. (2.5), as easily checked. This does not satisfy one of the conditions of the claim mentioned above. Thus, energy can be transferred from the gauge field into the charged scalar field, like particle creation by an external field.

In the asymptotically flat case, the superradiance scattering by a complex scalar field occurs for RN black holes and the reflected wave is amplified by the scattering potential [18]. The condition for its occurrence depends on the sign of the gauge coupling q . In this sense, the instabilities found in Sec. II is different from this usual picture of superradiance since ω_I in Eq. (2.20b) does not depend on the sign of q . A detailed analysis would be necessary in this direction.

In this article, we restrict our interest into the pure-electric case. As shown in Ref. [14], the plane-symmetric RN-AdS₄ possesses an inhomogeneous scalar hair with a vortex lattice structure under the probe approximation when magnetic field is included. It is one of interesting directions of our research to investigate what happens when the back-reaction of the inhomogeneous scalar field onto the geometry.

Acknowledgments

We would like to thank Makoto Natsuume, Takashi Okamura, Hideo Kodama for useful discussions. This research was supported in part by the Grant-in-Aid for Scientific Research (20540285) from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

Appendix A: The first law of black hole thermodynamics

In this appendix, we verify that the first law of thermodynamics holds for the asymptotically anti-de Sitter black hole with the complex scalar hair ψ , for any value of k of the 2-dimensional Einstein space. To do so, we follow the method developed by Wald and collaborators [19]. Although it can be applicable to rotating black holes, for simplicity, we focus on the static black holes with the 2-dimensional Einstein space of constant curvature.

We start with the variation of $\varepsilon_{c_1 c_2 c_3 c_4} \mathcal{L}(g_{ab}, A_a, \psi)$, which is written as

$$\delta [\varepsilon_{c_1 c_2 c_3 c_4} \mathcal{L}(\mathcal{F})] = \varepsilon_{c_1 c_2 c_3 c_4} \nabla_b \Theta^b(\mathcal{F}, \delta\mathcal{F}), \quad (\text{A1})$$

where $\varepsilon_{c_1 c_2 c_3 c_4}$ is the volume element of the spacetime, $\mathcal{F}$ denotes the field variables g_{ab} , A_a , and ψ , collectively. $\Theta^b(\mathcal{F}, \delta\mathcal{F})$ in Eq. (A1) is derived for the Lagrangian density (2.1) as

$$\Theta^b(\mathcal{F}, \delta\mathcal{F}) = \Theta_{(g)}^b(\mathcal{F}, \delta\mathcal{F}) + \Theta_{(A)}^b(\mathcal{F}, \delta\mathcal{F}) + \Theta_{(\psi)}^b(\mathcal{F}, \delta\mathcal{F}), \quad (\text{A2})$$

where

$$\Theta_{(g)}^b(\mathcal{F}, \delta\mathcal{F}) = g_{cd} \nabla^b \delta g^{cd} - \nabla_c \delta g^{bc}, \quad (\text{A3})$$

$$\Theta_{(A)}^b(\mathcal{F}, \delta\mathcal{F}) = -F^{bc} \delta A_c, \quad (\text{A4})$$

$$\Theta_{(\psi)}^b(\mathcal{F}, \delta\mathcal{F}) = -(D^b \psi) \delta \psi^\dagger - (D^b \psi)^\dagger \delta \psi. \quad (\text{A5})$$

When the variation δ is given by infinitesimal diffeomorphism, i.e., the Lie derivative L_η along an arbitrary vector η^a , the current $J^b(\mathcal{F}, \eta)$ defined by

$$J^b(\mathcal{F}, \eta) \equiv \Theta^b(\mathcal{F}, L_\eta \mathcal{F}) - \eta^b \mathcal{L}(\mathcal{F}) \quad (\text{A6})$$

is conserved on shell, and we can find the Noether potential $Q^{ba}(\mathcal{F}, \eta)$ of $J^b(\mathcal{F}, \eta)$, such that

$$J^b(\mathcal{F}, \eta) \equiv \nabla_a Q^{ba}(\mathcal{F}, \eta). \quad (\text{A7})$$

For the Lagrangian density (2.1), $Q^{ba}(\mathcal{F}, \eta)$ is derived as

$$Q^{ba}(\mathcal{F}, \eta) = Q_{(g)}^{ba}(\mathcal{F}, \eta) + Q_{(A)}^{ba}(\mathcal{F}, \eta), \quad (\text{A8})$$

where

$$Q_{(g)}^{ba}(\mathcal{F}, \eta) = \nabla^a \eta^b - \nabla^b \eta^a, \quad (\text{A9})$$

$$Q_{(A)}^{ba}(\mathcal{F}, \eta) = F^{ab} A_c \eta^c, \quad (\text{A10})$$

and the contribution to $Q^{ba}(\mathcal{F}, \eta)$ from the scalar field is absent [20].

In particular, when η^a is a Killing vector ξ^a , one can show

$$0 = \delta \int_{\partial\Sigma} \frac{1}{2} \varepsilon_{abcd} Q^{ba}(\mathcal{F}, \xi) + \int_{\partial\Sigma} \varepsilon_{abcd} \xi^b \Theta^a(\mathcal{F}, \delta\mathcal{F}), \quad (\text{A11})$$

for an arbitrary hypersurface Σ . In order to derive the first law, we choose as ξ^a the timelike Killing vector k^a

that becomes null at the horizon, and Σ to be a space-like hypersurface connecting the spacial infinity and the bifurcation surface $\mathcal{B}$ of the black hole horizon, which we assume to exist.

Thus, the boundary $\partial\Sigma$ includes the whole (in the case of $k = 1$) or a portion (for $k = 0$ and $k = -1$) of $\mathcal{B}$ and the 2-dimensional Einstein space at infinity. In addition, in the case of $k = 0$ or $k = -1$, there are other parts of the boundary, because the 2-dimensional Einstein space is not closed. However, in this case, we respect the isometries of the 2-dimensional Einstein space in choosing Σ , and impose a periodic boundary condition at these parts of the boundary. Then, the contribution from these parts of the boundary vanishes in Eq. (A11).

We define the total mass (for $k = 1$) or the mass density (for $k = 0$ and $k = -1$) of the black hole, which we denote as M , by

$$M \equiv \frac{1}{4w_2} \int_{\infty} \frac{1}{2} \varepsilon_{abcd} \left[Q_{(g)}^{ba}(\mathcal{F}, k) + 2 k^{[b} B^{a]}(\mathcal{F}) \right], \quad (\text{A12})$$

where $B^a(\mathcal{F})$ is defined by

$$\delta \int_{\infty} \varepsilon_{abcd} k^b B^a(\mathcal{F}) = \int_{\infty} \varepsilon_{abcd} k^b \Theta_{(g)}^a(\mathcal{F}, \delta\mathcal{F}), \quad (\text{A13})$$

and w_2 is the area of the whole (for $k = 1$) or the portion (for $k = 0$ and $k = -1$) of the 2-dimensional Einstein space with the metric $d\Omega_2^2$. It has been shown [21] that the mass defined by Eq. (A12) coincides with other various definitions of mass. The electric charge (or charge density) ρ of the black hole is defined as

$$\rho \equiv \frac{1}{w_2} \int_{\infty} \frac{1}{2} \varepsilon_{abcd} F^{ba}, \quad (\text{A14})$$

and the static potential is given by $-k^a A_a$, as usual. The gauge invariant quantity associated with the static potential, in the first law of black hole thermodynamics, is the difference of its value between the horizon and infinity. Since $k^a = 0$ on $\mathcal{B}$, this quantity is shown, in a regular gauge, to coincide with the chemical potential μ of the field theory, due to AdS/CFT, as

$$(-k^a A_a|_{\mathcal{B}}) - (-k^a A_a|_{\infty}) = k^a A_a|_{\infty} = \mu. \quad (\text{A15})$$

Furthermore, for the boundary condition Eq. (2.11) of the scalar field, $\psi \sim r^{-\Delta_+}$, and we have

$$\int_{\infty} \varepsilon_{abcd} k^b \Theta_{(\psi)}^a(\mathcal{F}, \delta\mathcal{F}) = O(r^{-2\Delta_++3}), \quad (\text{A16})$$

which shows that the contribution of the scalar field at infinity vanishes since $\Delta_+ > 3/2$.

Therefore, by defining the entropy (or entropy density) S of the black hole as

$$TS \equiv \frac{1}{4w_2} \int_{\mathcal{B}} \frac{1}{2} \varepsilon_{abcd} Q^{ba}(\mathcal{F}, k), \quad (\text{A17})$$

and noting $k^{[a}F^{bc]} = 0$ and the fact that k^a vanishes on $\mathcal{B}$, we obtain the first law as

$$T\delta S = \delta M - \frac{\mu}{4}\delta\rho. \quad (\text{A18})$$

While one can explicitly derive Eq. (A18) for the RN-AdS₄ black hole, from Eqs. (2.3c), (2.4), and (2.5), it is satisfied also for hairy black holes.

-
- [1] S. A. Hartnoll, “Lectures on holographic methods for condensed matter physics,” *Class. Quant. Grav.* **26**, 224002 (2009) [arXiv:0903.3246 [hep-th]].
 - [2] C. P. Herzog, “Lectures on holographic superfluidity and superconductivity,” *J. Phys. A* **42**, 343001 (2009) [arXiv:0904.1975 [hep-th]].
 - [3] G. T. Horowitz, “Introduction to Holographic Superconductors,” arXiv:1002.1722[hep-th].
 - [4] J. M. Maldacena, “The large N limit of superconformal field theories and supergravity,” *Adv. Theor. Math. Phys.* **2** (1998) 231 [*Int. J. Theor. Phys.* **38** (1999) 1113] [arXiv:hep-th/9711200].
 - [5] S. A. Hartnoll, C. P. Herzog, and G. T. Horowitz, “Holographic superconductors,” *JHEP* **0812**, (2008) 015 [arXiv:0810.1563 [hep-th]].
 - [6] P. Breitenlohner and D. Z. Freedman, “Stability In Gauged Extended Supergravity,” *Annals Phys.* **144** (1982) 249; “Positive Energy In Anti-De Sitter Backgrounds And Gauged Extended Supergravity,” *Phys. Lett. B* **115** (1982) 197.
 - [7] T. Torii, K. Maeda, and M. Narita, “Scalar hair on the black hole in asymptotically anti-de Sitter spacetime,” *Phys. Rev. D* **64** 044007 (2001).
 - [8] T. Hertog and K. Maeda, “Black Holes with Scalar Hair and Asymptotics in N=8 Supergravity,” *JHEP* **0407**, (2004) 051 [arXiv:hep-th/0404261].
 - [9] S. S. Gubser, “Breaking an Abelian gauge symmetry near a black hole horizon,” *Phys. Rev. D* **78**, 065034 (2008) [arXiv: 0801.2977 [hep-th]].
 - [10] K. Maeda, M. Natsuume, and T. Okamura, “Dynamic critical phenomena in the AdS/CFT duality,” *Phys. Rev. D* **78**, 106007 (2008) [arXiv:0809.4074 [hep-th]].
 - [11] K. Maeda, M. Natsuume, and T. Okamura, “Universality class of holographic superconductors,” *Phys. Rev. D* **79**, 126004 (2009) [arXiv:0904.1914 [hep-th]].
 - [12] T. Hertog and G. T. Horowitz, “Towards a big crunch dual,” *JHEP* **0407**, (2004) 073 [arXiv:hep-th/0406134]; T. Hertog and G. T. Horowitz, “Holographic description of AdS cosmologies,” *JHEP* **0504**, (2005) 005 [arXiv:hep-th/0503071].
 - [13] K. Maeda and T. Okamura, “Characteristic length of an AdS/CFT superconductor,” *Phys. Rev. D* **78**, 106006 (2008) [arXiv:0809.3079 [hep-th]].
 - [14] K. Maeda, M. Natsuume, and T. Okamura, “Vortex lattice for a holographic superconductor,” *Phys. Rev. D* **81**, 026002 (2010) [arXiv:0910.4475 [hep-th]].
 - [15] V. Cardoso and O. J. C. Dias, *Phys. Rev. D* **70**, 084011 (2004) [arXiv:hep-th/0405006]; V. Cardoso, O. J. C. Dias, J. P. S. Lemos, and S. Yoshida, *Phys. Rev. D* **70**, 044039 (2004) (Erratum: *ibid.* **D70**:049903 (2004)) [arXiv:hep-th/0404096]; V. Cardoso, O. J. C. Dias, and S. Yoshida, *Phys. Rev. D* **74**, 044008 (2006) [arXiv:hep-th/0607162].
 - [16] S. Hawking and H. Reall, “Charged and rotating AdS black holes and their CFT duals,” *Phys. Rev. D* **61**, 024014 (1999) [arXiv:hep-th/9908109].
 - [17] H. Kodama, “Superradiance and Instability of Black Holes,” *Prog. Theor. Phys. Suppl.* **172**, 11 (2008).
 - [18] G. W. Gibbons, *Commun. Math. Phys.* **44** 245 (1975).
 - [19] R. M. Wald, *Phys. Rev. D* **48**, R3427 (1993), V. Iyer and R.M. Wald, *Phys. Rev. D* **50**, 846 (1994).
 - [20] T. Jacobson, G. Kang, and R. C. Myers, *Phys. Rev. D* **49** 6587 (1994).
 - [21] S. Hollands, A. Ishibashi, and D. Marolf, *Class. Quant. Grav.* **22** 2881 (2005).